

Repositório ISCTE-IUL

Deposited in *Repositório ISCTE-IUL*:

2026-04-10

Deposited version:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Mendes, S., Soares, H. & Miró-Roig, M. (2021). Vector bundles E on P^3 with homological dimension 2 and $\chi(\text{End } E)=1$. *Forum Mathematicum*. 33 (3), 808-820

Further information on publisher's website:

[10.1515/forum-2020-0169](https://doi.org/10.1515/forum-2020-0169)

Publisher's copyright statement:

This is the peer reviewed version of the following article: Mendes, S., Soares, H. & Miró-Roig, M. (2021). Vector bundles E on P^3 with homological dimension 2 and $\chi(\text{End } E)=1$. *Forum Mathematicum*. 33 (3), 808-820, which has been published in final form at <https://dx.doi.org/10.1515/forum-2020-0169>. This article may be used for non-commercial purposes in accordance with the Publisher's Terms and Conditions for self-archiving.

Use policy

Creative Commons CC BY 4.0

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a link is made to the metadata record in the Repository
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

On the existence of vector bundles E of homological dimension 2 and $\chi(\text{End } E) = 1$

Sérgio Mendes, Rosa María Miró-Roig and Helena Soares

Abstract

We find the complete integer solutions of the equation

$$X^2 + Y^2 + Z^2 - 4XY - 4YZ + 10XZ = 1.$$

As an application we prove that for each solution (a, b, c) such that $a > 0$, $b - 2a > 0$ and $(b - 2a)^2 \geq 4a$ there is a vector bundle E on \mathbb{P}^3 defined by a minimal linear resolution

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2)^a \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^b \rightarrow \mathcal{O}_{\mathbb{P}^3}^c \rightarrow E \rightarrow 0.$$

In particular, E satisfies $\chi(\text{End } E) = 1$.

1 Introduction

Let E be a vector bundle on \mathbb{P}^n , with $n \geq 2$, of homological dimension 1 and defined by a minimal linear resolution

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^n}(-1)^a \rightarrow \mathcal{O}_{\mathbb{P}^n}^b \rightarrow E \rightarrow 0. \quad (1)$$

These bundles were first introduced by Dolgachev and Kapranov [5] and they are usually called Steiner bundles. Their study is extensive and has been addressed from very different points of view. We refer in particular to [2], [3], [9], [10], [11].

As it can be easily deduced, e.g. from the formula given in Proposition 3.4 in [11], the Euler characteristic of $\text{End}(E)$ is

$$\chi(\text{End}(E)) = a^2 + b^2 - (n + 1)ab.$$

It is worth recalling that often one is interested in studying vector bundles satisfying $\chi(\text{End } E) = 1$ for they provide good candidates for exceptional

bundles, important objects when studying stability and derived categories. We know that for $n \geq 2$ all positive integer solutions of the quadratic form

$$X^2 + Y^2 - (n + 1)XY = 1 \quad (2)$$

are pairs $(X, Y) = (u_k, u_{k+1})$, with $k \geq 1$, where $\{u_k\}_{k \geq 0}$ is the sequence defined recursively by

$$\begin{cases} u_0 = 0 \\ u_1 = 1 \\ u_{k+2} = (n + 1)u_{k+1} - u_k \end{cases}$$

So, if E is a vector bundle as in (1) and $\chi(\text{End}(E)) = 1$ then $a = u_k$ and $b = u_{k+1}$, for some $k \geq 1$. Conversely, for each solution pair (u_k, u_{k+1}) of the equation (2) we know that there exists a vector bundle E defined by an exact sequence of type (1), with $a = u_k$ and $b = u_{k+1}$. In fact, in this case we know how to construct all such vector bundles (see Lemma 2.2.5 and Theorem 2.2.7 in [13]).

In the present work we address the case of homological dimension 2. More precisely, we consider a non-splitting vector bundle E defined by a minimal linear resolution

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^n}(-2)^a \rightarrow \mathcal{O}_{\mathbb{P}^n}(-1)^b \rightarrow \mathcal{O}_{\mathbb{P}^n}^c \rightarrow E \rightarrow 0, \quad (3)$$

with $n \geq 3$. From Proposition 3.4 in [11] the Euler characteristic of $\text{End } E$ is the ternary quadratic form

$$\chi(\text{End}(E)) = a^2 + b^2 + c^2 - (n + 1)ab - (n + 1)bc + \binom{n + 2}{2}ac.$$

If $\chi(\text{End}(E)) = 1$ then a , b and c are positive integer solutions of the Diophantine equation

$$X^2 + Y^2 + Z^2 - (n + 1)XY - (n + 1)YZ + \binom{n + 2}{2}XZ = 1. \quad (4)$$

We want to know if, given an arbitrary positive integer solution (a, b, c) of the above equation, there exists a vector bundle E with a linear resolution of type (3) and exponents (also called the Betti numbers) a , b and c .

The first obstacle we undergo is to find all positive integer solutions of (4). In section 2 we delve into the theory of ternary quadratic forms, following [4]. We begin by recalling some generalities and proceed with the classification of the family of ternary quadratic forms (4), for n up to 20, using Legendre's

theorem. Describing all positive solutions is a much harder problem than the corresponding one in the case of homological dimension 1, so we will need to restrict our study to $n = 3$ for which the above equation becomes

$$X^2 + Y^2 + Z^2 - 4XY - 4YZ + 10XZ = 1. \quad (5)$$

We end section 2 with the complete integers solutions of (5) (Proposition 2.6).

In section 3 we treat the existence problem and prove our main result, Theorem 3.3. For each solution (a, b, c) of equation (5) such that $a > 0$, $b - 2a > 0$ and $(b - 2a)^2 \geq 4a$, we construct a vector bundle E on \mathbb{P}^3 defined by a minimal linear resolution of length 2, thus satisfying $\chi(\text{End}(E)) = 1$. The main result behind is the existence of a curve C on \mathbb{P}^3 whose ideal sheaf \mathcal{I}_C is defined by a linear resolution of length 2 and such that $\text{Ext}^1(\mathcal{I}_C(b - 2a), \mathcal{O}_{\mathbb{P}^3}) \neq 0$ (see Proposition 3.2). Then we can take a non-trivial extension E of $\mathcal{O}_{\mathbb{P}^3}$ by $\mathcal{I}_C(b - 2a)$ and prove that E has the required resolution.

In [11] the authors approached the problem of characterising exceptional vector bundles of arbitrary homological dimension. They were able to explicitly construct a family of exceptional vector bundles of homological dimension 2 and linear resolution on the projective space \mathbb{P}^n (Proposition 4.3). When $n = 3$ this family is a subset of the set of vector bundles constructed herein.

Notation 1.1. *If E is a vector bundle on \mathbb{P}^n we denote $h^q(E) = \dim H^q(E) = \dim H^q(\mathbb{P}^n, E)$. The Euler characteristic of E is defined by the integer*

$$\chi(E) = \sum_{i=0}^n (-1)^i h^i(E).$$

Recall furthermore that we have an isomorphism $\text{End } E \cong E^\vee \otimes E$.

The minimal length of a resolution of E , i.e. the homological dimension of E , is denoted by $\text{hd}(E)$.

2 The solutions of the Diophantine equation

$$\chi(\text{End } E) = 1$$

Before we specify to the case $n = 3$ let us consider the more general family of Diophantine equations (4), arising from the algebraic geometric problem described in the introduction

$$X^2 + Y^2 + Z^2 - (n + 1)XY - (n + 1)YZ + \binom{n + 2}{2}XZ = 1, \quad n \geq 3,$$

where $\binom{n}{k}$ denote the binomial coefficients.

The solution of this Diophantine equation is related to the classical problem of representing integers by ternary quadratic forms. A well-known example, solved by Gauss [12, p. 79], is to determine which integers can be written in the form $x^2 + y^2 + z^2$.

The first step is to write the above equation in a reduced form. Then, using Legendre's theorem [8], we classify each reduced form for a given $n \geq 3$ as being either isotropic or anisotropic. Each one of these two cases has a different technique. As mentioned before, we shall concentrate in the isotropic case $n = 3$ using the methods described in [4]. As we will see, although the method is explained in Cassel's book, explicit formulae of the integral solutions for a given n are not easy to obtain. In Cassel's own words [4, p. 303]: "The numerical details could be tedious in any given case."

2.1 The reduced equations

Equation (4) can be written as

$$\left(\frac{n+1}{2}X - Y + \frac{n+1}{2}Z\right)^2 - \frac{n^2+3n-2}{8}(X-Z)^2 - \frac{n^2+n-4}{8}(X+Z)^2 = 1$$

and also

$$((n+1)X - 2Y + (n+1)Z)^2 - \frac{n^2+n-4}{2}(X+Z)^2 - \frac{n^2+3n-2}{2}(X-Z)^2 = 4.$$

Write

$$a = a_n = \frac{n^2+n-4}{2} \text{ and } b = b_n = \frac{n^2+3n-2}{2}. \quad (6)$$

Observe that a_n and b_n are positive integers for $n \geq 2$ (recall that we are concerned only with $n \geq 3$). Moreover, a_n and b_n are related by

$$b_n = a_n + n + 1, \forall n.$$

Now, we make a first change of variables

$$u = (n+1)X - 2Y + (n+1)Z, \quad v = X + Z, \quad w = X - Z.$$

Equation (4) becomes

$$u^2 = 4 + \frac{n^2+n-4}{2}v^2 + \frac{n^2+3n-2}{2}w^2,$$

or with the notation introduced by (6), simply

$$u^2 = 4 + av^2 + bw^2.$$

Rewriting this last equation as

$$\left(\frac{u}{2}\right)^2 - a\left(\frac{v}{2}\right)^2 - b\left(\frac{w}{2}\right)^2 = 1$$

and performing a second change of variables

$$x = \frac{u}{2}, \quad y = \frac{v}{2}, \quad z = \frac{w}{2},$$

we obtain

$$x^2 - ay^2 - bz^2 = 1. \tag{7}$$

If a and b are not perfect squares then $f = x^2 - ay^2 - bz^2$ is the reduced ternary form associated with equation (4). Otherwise, we need to change variables one last time to obtain the reduced ternary form.

2.2 Notation and terminology

For the sake of completeness we recall some basic terminology about rational quadratic forms (see [4] for a detailed account). As usual \mathbb{Q} denotes the field of rational numbers and \mathbb{Z} the ring of integers. Let \mathcal{I} be a ring in \mathbb{Q} . We allow the possibility $\mathcal{I} = \mathbb{Q}$. We say that a quadratic form

$$f(x_1, \dots, x_n) = \sum_{i,j} a_{ij}x_ix_j,$$

with $a_{ij} = a_{ji} \in \mathbb{Q}$ in n variables x_1, \dots, x_n , represents an element $c \in \mathbb{Q}$ over \mathcal{I} if there is some $\mathbf{b} = (b_1, \dots, b_n) \in \mathcal{I}^n$ such that $f(\mathbf{b}) = c$. In particular, we shall speak of rational representations if $\mathcal{I} = \mathbb{Q}$ and of integral representations if $\mathcal{I} = \mathbb{Z}$. In this note we are only concerned with integral representations. The quadratic form is integer-valued if $f(\mathbf{b}) \in \mathbb{Z}$, for all $\mathbf{b} \in \mathcal{I}^n$. In terms of coefficients it means that

$$a_{ii} \in \mathcal{I} \text{ and } 2a_{ij} \in \mathcal{I} \text{ (} 1 \leq i \leq j \leq n \text{)}.$$

We say that f is classically integral if

$$a_{ij} \in \mathcal{I} \text{ (} 1 \leq i \leq n, 1 \leq j \leq n \text{)},$$

that is, if the elements of the Gram matrix are in \mathcal{I} . When f is classically integral we say that it is primitive if $\text{g.c.d.}(a_{ij}) = 1$, where g.c.d. denotes the greatest common divisor.

A quadratic form f is said to be isotropic if it represents 0 nontrivially, otherwise it is anisotropic.

Two forms f and g are \mathbb{Z} -equivalent if there is a square integral matrix T such that

$$g(\mathbf{X}) = f(T\mathbf{X})$$

and $\det(T) \in \mathcal{U}(\mathbb{Z}) = \{\pm 1\}$, where $\mathcal{U}(\mathbb{Z})$ is the group of units of \mathbb{Z} . The equivalence is proper if $\det(T) = +1$ and improper if $\det(T) = -1$. Clearly, proper equivalence is an equivalence relation. An equivalence of a form f with itself is an integral automorph. Denote by $\mathcal{O}(f)$ the set of automorphs of f and by \mathcal{O}^+ the set of proper automorphs. $\mathcal{O}^+(f)$ forms a group under composition, the group of (proper) automorphs of f , also called the orthogonal group of f .

We also need some terminology on binary quadratic forms. Let

$$f = ax^2 + bxy + cy^2$$

be an integral binary quadratic form in the non-classical definition. The discriminant of f is given by $\Delta = b^2 - 4ac$, and it is related with the determinant $d(f)$ of f , i.e. of the associated Gram matrix, by $\Delta(b) = -4d(f)$.

2.3 Classification of ternary forms

In order to decide if an indefinite ternary quadratic form is isotropic or anisotropic we may use the beautiful theorem of Legendre (see [14, Ch. II, §XIV, Ch. IV and Appendix I] for a historical perspective). The version bellow can be found in [8, Proposition 17.3.2] and is more suitable for our purposes. Let \mathcal{R} denote the following binary relation over the set of nonzero integers: given $m, n \in \mathbb{Z} \setminus \{0\}$,

$$m\mathcal{R}n \Leftrightarrow x^2 = m \pmod{n},$$

for some integer x .

Theorem 2.1 (Legendre's theorem). *Let a, b be positive squarefree integers. Then,*

$$x^2 = ay^2 + bz^2$$

has a nontrivial solution, that is, the form $f = x^2 - ay^2 - bz^2$ is isotropic if, and only if, the following conditions are satisfied:

- (i) $a\mathcal{R}b$;
- (ii) $b\mathcal{R}a$;
- (iii) $-\frac{ab}{d^2}\mathcal{R}d$, where $d = \text{g.c.d.}(a, b)$.

The following remark is useful for explicit computations.

Remark 2.2. Note that if b is odd, then

$$a\mathcal{R}b \Rightarrow \left(\frac{a}{b}\right) = 1,$$

where (\div) denotes the Jacobi symbol. We recall that, when $b = p$ is prime, the Legendre symbol is defined to be

$$\left(\frac{a}{p}\right) = \begin{cases} 1 & , \text{ if } x^2 \equiv a \pmod{p} \text{ has a nonzero solution} \\ -1 & , \text{ if } x^2 \equiv a \pmod{p} \text{ has no solution,} \end{cases}$$

and $\left(\frac{a}{p}\right) = 0$ when p divides a . The Jacobi symbol generalizes the Legendre symbol as follows. Given an arbitrary odd integer b , if $b = p_1^{e_1} \dots p_k^{e_k}$, where p_i are odd primes, set

$$\left(\frac{a}{b}\right) = \left(\frac{a}{p_1}\right)^{e_1} \dots \left(\frac{a}{p_k}\right)^{e_k}.$$

Example 2.3. For $n = 4$, the (reduced) ternary form (7) is

$$x^2 = 8y^2 + 13z^2 = 2(2y)^2 + 13z^2.$$

With the obvious change of variables we may represent it by

$$x^2 = 2y^2 + 13z^2.$$

Now, $13\mathcal{R}2$ since $9 = 13 - 2 \times 2$. However,

$$\left(\frac{2}{13}\right) = -\left(\frac{1}{13}\right) = -1$$

and so, by the above remark, the form is anisotropic.

We now present a list of reduced ternary quadratic forms associated with equation (7) for n from 3 up to 20. The classification of each form $f = x^2 - ay^2 - bz^2$ is obtained using Legendre's theorem and some basic computations with the Jacobi symbol.

n	$f = x^2 - ay^2 - bz^2$	<i>Classification</i>
3	$x^2 - 4y^2 - 8z^2$	<i>isotropic</i>
4	$x^2 - 8y^2 - 13z^2$	<i>anisotropic</i>
5	$x^2 - 13y^2 - 19z^2$	<i>anisotropic</i>
6	$x^2 - 19y^2 - 29z^2$	<i>anisotropic</i>
7	$x^2 - 26y^2 - 34z^2$	<i>anisotropic</i>
8	$x^2 - 34y^2 - 43z^2$	<i>anisotropic</i>
9	$x^2 - 43y^2 - 53z^2$	<i>isotropic</i>
10	$x^2 - 54y^2 - 64z^2$	<i>isotropic</i>
11	$x^2 - 64y^2 - 76z^2$	<i>isotropic</i>
12	$x^2 - 76y^2 - 89z^2$	<i>anisotropic</i>
13	$x^2 - 89y^2 - 103z^2$	<i>anisotropic</i>
14	$x^2 - 103y^2 - 118z^2$	<i>anisotropic</i>
15	$x^2 - 118y^2 - 134z^2$	<i>anisotropic</i>
16	$x^2 - 134y^2 - 151z^2$	<i>anisotropic</i>
17	$x^2 - 151y^2 - 169z^2$	<i>isotropic</i>
18	$x^2 - 169y^2 - 188z^2$	<i>isotropic</i>
19	$x^2 - 188y^2 - 208z^2$	<i>anisotropic</i>
20	$x^2 - 208y^2 - 229z^2$	<i>anisotropic</i>

Table 1.

One case is particularly easy. In fact, the form is isotropic if either a or b is a square. For, suppose a is a square. Then, $(\sqrt{a}, 1, 0)$ is a nontrivial integral representation of 0 by f . This is the case when $n = 3, 10, 11, 17$ and 18, see table 1.

Example 2.4. Consider $n = 9$. The form is $f = x^2 - 43y^2 - 53z^2$ and both 43 and 53 are squarefree (actually, prime numbers). We have

$$19^2 = 43 + 6 \times 53$$

so $43\mathcal{R}53$, and also

$$15^2 = 53 + 4 \times 43$$

that is, $53\mathcal{R}43$. By Legendre's theorem, f is isotropic.

2.4 Isotropic ternary forms

It turns out that the study of the representation of integers by isotropic ternary forms reduces to the study of the following form

$$f_0(x, y, z) = xz - y^2.$$

We now follow [4, p. 301]. Let

$$S = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in SL^\pm(2, \mathbb{Z}),$$

where $SL^\pm(2, \mathbb{Z})$ is the modular group of elements of $GL(2, \mathbb{Z})$ with determinant ± 1 . Then,

$$T(S) = \tilde{S} = \begin{pmatrix} \alpha^2 & 2\alpha\gamma & \gamma^2 \\ \alpha\beta & \alpha\delta + \beta\gamma & \gamma\delta \\ \beta^2 & 2\beta\gamma & \delta^2 \end{pmatrix}$$

is an automorph of f_0 .

According to [4, Lemma 5.2] the map $S \mapsto T(S)$ is a homomorphism of $SL^\pm(2, \mathbb{Z})$ onto $\mathcal{O}^+(f_0)$ with kernel $\{\pm I\}$. In other words, we have the following description of the group of automorphs of f_0 :

$$\mathcal{O}^+(f_0) \cong SL^\pm(2, \mathbb{Z})/\{\pm I\}.$$

Given any other isotropic ternary form f , there exist an integer m and an integral matrix M such that

$$mf(\mathbf{X}) = f_0(M\mathbf{X}),$$

see [4, p. 303]. Hence, any integral representation $\mathbf{b} \in \mathbb{Z}^3$ of a given integer $r \in \mathbb{Z}$ by f gives rise to the integral representation $M\mathbf{b}$ of mr by f_0 . On the other hand, every representation of mr by f_0 is of the type $T(S)c$, where c runs through a finite set, say Φ_f , and $S \in SL^\pm(2, \mathbb{Z})/\{\pm I\}$. Furthermore, $T(S)c$ comes from an integral representation of r by f whenever

$$M^{-1}T(S)c \in \mathbb{Z}^3. \tag{8}$$

We conclude that in order to find all the integral solutions of the Diophantine equation (7),

$$x^2 - ay^2 - bz^2 = 1,$$

with $f = x^2 - ay^2 - bz^2$ isotropic, it amounts to:

- (i) Find M and m ;
- (ii) Identify the finite set of $\Phi_f = \{c_1, \dots, c_k\}$;
- (iii) Solve a system of three modular equations (*mod* $\det(M)$).

If a and b in equation (7) are both squarefree, it can be difficult to determine explicitly M and m . These computations are not explained in Cassel's book. But the case where a or b is a square is fairly simple.

Proposition 2.5. *Let $f = x^2 - ay^2 - bz^2$ with a and b positive integers. We have:*

(i) *If a is a square then f is isotropic and we may take*

$$m = \tilde{b} \quad \text{and} \quad M = M_{\tilde{b}} = \begin{bmatrix} \tilde{b} & \tilde{b} & 0 \\ 0 & 0 & \tilde{b} \\ 1 & -1 & 0 \end{bmatrix},$$

where $\tilde{b}|b$ and \tilde{b} has squarefree prime factors.

(ii) *If b is a square then f is isotropic and we may take*

$$m = \tilde{a} \quad \text{and} \quad M = M_{\tilde{a}} = \begin{bmatrix} \tilde{a} & 0 & \tilde{a} \\ 0 & \tilde{a} & 0 \\ 1 & 0 & -1 \end{bmatrix},$$

where $\tilde{a}|a$ and \tilde{a} has squarefree prime factors.

Proof. (i) Since a is a square we obtain a new ternary form, which is primitive,

$$\tilde{f} = x^2 - (\sqrt{a}y)^2 - \tilde{b}(\sqrt{c}z)^2,$$

with the obvious change of variables. Here, $b = \tilde{b}c$, \tilde{b} is a squarefree factor and c is a square (with the possibility that $\sqrt{c} = 1$). In matrix form, $f_0(M_{\tilde{b}}\mathbf{X})$ with $M_{\tilde{b}}$ as above becomes:

$$\begin{aligned} M_{\tilde{b}}^t M_{f_0} M_{\tilde{b}} &= \begin{bmatrix} \tilde{b} & 0 & 1 \\ \tilde{b} & 0 & -1 \\ 0 & \tilde{b} & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1/2 \\ 0 & -1 & 0 \\ 1/2 & 0 & 0 \end{bmatrix} \begin{bmatrix} \tilde{b} & \tilde{b} & 0 \\ 0 & 0 & \tilde{b} \\ 1 & -1 & 0 \end{bmatrix} \\ &= \begin{bmatrix} \tilde{b} & 0 & 0 \\ 0 & -\tilde{b} & 0 \\ 0 & 0 & -\tilde{b}^2 \end{bmatrix} \\ &= \tilde{b} M_{\tilde{f}}. \end{aligned}$$

The case (ii) is similar. □

Now, we identify the set Φ_f . By [4, Lemma 5.3], the elements c_i correspond to representatives of proper equivalence classes of classically integral binary forms of determinant m . Surely, there are only finitely many such classes.

2.5 The case $n = 3$

Now, we specify to the case $n = 3$. The quadratic form $f = x^2 - 4y^2 - 8z^2$ is clearly isotropic, for $(\pm 2, \pm 1, 0)$ are nontrivial representations of 0 by f . We want to find all the integral solutions of

$$f = 1 \Leftrightarrow x^2 - (2y)^2 - 2(2z)^2 = 1. \quad (9)$$

With the change variables

$$X_1 = x, \quad X_2 = 2y, \quad X_3 = 2z$$

we have

$$f(X_1, X_2, X_3) = X_1^2 - X_2^2 - 2X_3^2 = 1. \quad (10)$$

Equation (10) is the reduced, primitive form of equation (5). From the above discussion, there exists $m \in \mathbb{Z}$ and an integral 3×3 matrix M such that

$$mf(\mathbf{X}) = f_0(M\mathbf{X}),$$

with $\mathbf{X} = (X_1, X_2, X_3) \in \mathbb{Z}^3$. By Proposition 2.5, we have $m = 2$ and

$$M = \begin{bmatrix} 2 & 2 & 0 \\ 0 & 0 & 2 \\ 1 & -1 & 0 \end{bmatrix}. \text{ Then}$$

$$f(\mathbf{X}) = 1 \Rightarrow f_0(M\mathbf{X}) = 2 \times 1 = 2.$$

There is only one proper equivalence class of classically integral binary forms of determinant 2: it corresponds to the reduced binary form $(1, 0, 2)$. It has, of course, discriminant $\Delta = -8$.

Now, from (8), the solutions of equation (10) are given by

$$\begin{aligned} M^{-1}T(S)c &= \frac{1}{8} \begin{bmatrix} 2 & 0 & 4 \\ 2 & 0 & -4 \\ 0 & 4 & 0 \end{bmatrix} T(S)c \\ &= \frac{1}{4} \begin{bmatrix} 1 & 0 & 2 \\ 1 & 0 & -2 \\ 0 & 2 & 0 \end{bmatrix} \begin{bmatrix} \alpha^2 & 2\alpha\gamma & \gamma^2 \\ \alpha\beta & \alpha\delta + \beta\gamma & \gamma\delta \\ \beta^2 & 2\beta\gamma & \delta^2 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} \\ &= \frac{1}{4} \begin{bmatrix} \alpha^2 + 2\gamma^2 + 2\beta^2 + 4\delta^2 \\ \alpha^2 + 2\gamma^2 - 2\beta^2 - 4\delta^2 \\ 2\alpha\beta + 4\gamma\delta \end{bmatrix}. \end{aligned} \quad (11)$$

Therefore, we obtain a system with three modular equations:

$$\begin{cases} \alpha^2 + 2\gamma^2 + 2\beta^2 + 4\delta^2 \equiv 0 \pmod{4} \\ \alpha^2 + 2\gamma^2 - 2\beta^2 - 4\delta^2 \equiv 0 \pmod{4} \\ 2\alpha\beta + 4\gamma\delta \equiv 0 \pmod{4} \end{cases}$$

$$\begin{aligned} &\Leftrightarrow \begin{cases} \alpha^2 + 2\gamma^2 + 2\beta^2 \equiv 0 \pmod{4} \\ 2\alpha\beta \equiv 0 \pmod{4} \end{cases} \\ &\Rightarrow \begin{cases} \alpha^2 + 2\gamma^2 + 2\beta^2 \equiv 0 \pmod{2} \\ 2\alpha\beta \equiv 0 \pmod{2} \end{cases} \end{aligned}$$

We conclude that

$$\alpha^2 \equiv 0 \pmod{2}.$$

In other words, the integral solutions of the Diophantine equation (10) are parametrised by

$$\begin{bmatrix} \alpha & \beta \\ \gamma & \delta \end{bmatrix} \in SL^\pm(2, \mathbb{Z}),$$

with α even.

From the above, we may now obtain the integer solutions of equation (5) applying formulae (11), and then going backwards doing the proper change of variables. Hence, we have:

Proposition 2.6. *The integer solutions of equation (5) are given by*

$$\begin{aligned} X &= \frac{\alpha^2 + 2\gamma^2 - 2\beta^2 - 4\delta^2 + 2\alpha\beta + 4\gamma\delta}{8}, \\ Y &= \frac{\alpha^2 + 2\gamma^2 - 6\beta^2 - 12\delta^2}{4}, \\ Z &= \frac{\alpha^2 + 2\gamma^2 - 2\beta^2 - 4\delta^2 - 2\alpha\beta - 4\gamma\delta}{8}, \end{aligned} \tag{12}$$

with $\alpha, \beta, \gamma, \delta \in \mathbb{Z}$, α even and $\alpha\delta - \beta\gamma = \pm 1$.

3 Vector bundles on \mathbb{P}^3 of homological dimension 2 and $\chi(\text{End } E) = 1$

Let E be a vector bundle on \mathbb{P}^3 of homological dimension 2 defined by a minimal linear resolution

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2)^a \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^b \rightarrow \mathcal{O}_{\mathbb{P}^3}^c \rightarrow E \rightarrow 0, \tag{13}$$

and such that $\chi(\text{End } E) = 1$. Then, from Proposition 3.4 in [11], this is equivalent to the fact that the Betti numbers a , b and c satisfy

$$a^2 + b^2 + c^2 - 4ab - 4bc + 10ac = 1.$$

The 3-uple (a, b, c) is thus a solution of the Diophantine equation (5) and, as proved in Proposition 2.6, a, b and c are (positive) integers of the form (12), for some $\alpha, \beta, \gamma, \delta \in \mathbb{Z}$, α even and $\alpha\delta - \beta\gamma = \pm 1$.

Observe that according to this parameterisation we easily deduce that

$$\mathrm{rk} E = a - b + c = \beta^2 + 2\delta^2.$$

Now, we are interested in finding those integer solutions (12) which correspond to such a vector bundle E . It is obvious that there are infinitely many that do not give rise to any such bundle. First, we note that we are not able to find a complete description of all positive solutions of (5) as in the case of homological dimension 1. So for instance, if $\alpha = 0$ then $\beta = \pm 1$ and $\gamma = \pm 1$, and we get

$$a = \frac{-4\delta^2 \pm 4\delta}{8} = -\frac{1}{2}\delta^2 \pm \frac{1}{2}\delta \leq 0.$$

Second, the set of solutions obtained include cases we are excluding: in the case when $\delta = 0$, and so $\beta = \pm 1$, the rank of E is $\mathrm{rk} E = \beta^2 = 1$, contradicting the fact that $\mathrm{rk} E \geq 2$ (provided E is a non-splitting vector bundle, see [2], Corollary 1.7).

We next prove the following general fact concerning the first Chern class of any vector bundle defined by (13).

Proposition 3.1. *If E is a vector bundle on \mathbb{P}^3 of homological dimension 2 and minimal linear resolution*

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2)^a \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^b \rightarrow \mathcal{O}_{\mathbb{P}^3}^c \rightarrow E \rightarrow 0,$$

then it has positive first Chern class, i.e. $b - 2a > 0$.

Proof. Since E is globally generated with $h^0(E) = c$ and $r := \mathrm{rk} E \leq c$ we can take $r - 1$ global sections of E , $\sigma_1, \dots, \sigma_{r-1} \in H^0(E)$, and consider the curve

$$C = (\sigma_1 \wedge \dots \wedge \sigma_{r-1})_0,$$

defined as the degeneracy locus in \mathbb{P}^3 where $\sigma_1, \dots, \sigma_{r-1}$ are linearly dependent. We thus have a short exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}^{r-1} \rightarrow E \rightarrow \mathcal{I}_C(c_1(E)) \rightarrow 0,$$

where $c_1(E) = b - 2a$ denotes the first Chern class of E . Taking cohomology of this sequence we get

$$0 \rightarrow H^0(\mathcal{O}_{\mathbb{P}^3}^{r-1}) \cong \mathbb{C}^{r-1} \rightarrow H^0(E) \cong \mathbb{C}^c \rightarrow H^0(\mathcal{I}_C(c_1(E))) \rightarrow 0,$$

and so $H^0(\mathcal{I}_C(c_1(E))) \neq 0$. Hence C is contained in a surface of degree $c_1(E)$ and therefore, $c_1(E) > 0$. \square

In order to prove our main result, we will first prove the existence of curves in \mathbb{P}^3 whose ideal sheaf is defined by a minimal linear resolution of length 2.

For any positive integer p , denote

$$D_p := \frac{p(p+1)}{2}$$

and

$$d_p := \begin{cases} \frac{p(p+2)}{4}, & \text{if } p \text{ is even,} \\ \frac{(p+1)^2}{4}, & \text{if } p \text{ is odd.} \end{cases}$$

Proposition 3.2. *Suppose $p, s > 0$ are positive integers that satisfy*

$$\begin{cases} p^2 \geq 4s, & \text{if } p \text{ is even,} \\ p^2 - 1 \geq 4s, & \text{if } p \text{ is odd.} \end{cases} \quad (14)$$

Then there exists a smooth connected curve $C \subset \mathbb{P}^3$ with minimal resolution

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2-p)^s \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1-p)^{2s+p} \rightarrow \mathcal{O}_{\mathbb{P}^3}(-p)^{s+p+1} \rightarrow \mathcal{I}_C \rightarrow 0,$$

maximal rank, degree

$$\deg(C) = \binom{p+1}{2} - s,$$

and arithmetic genus

$$p_a = \frac{(2p+3)(p-1)(p-2)}{6} - s(p-1).$$

Furthermore, $H^2(\mathbb{P}^3, \mathcal{I}_C(p-4)) \neq 0$.

Proof. Let $d = \binom{p+1}{2} - s$ and let us first prove that $d_p \leq d \leq D_p$. In fact, on the one hand

$$d = \binom{p+1}{2} - s = \frac{(p+1)p}{2} - s \leq \frac{(p+1)p}{2} = D_p.$$

On the other hand, if p is even, we have

$$\begin{aligned} d_p &= \frac{p(p+2)}{4} \leq \binom{p+1}{2} - s = d \\ \Leftrightarrow \frac{p(p+2)}{4} &\leq \frac{(p+1)p}{2} - s \\ \Leftrightarrow 4s &\leq p^2. \end{aligned}$$

If p is odd, we similarly obtain

$$d_p = \frac{(p+1)^2}{4} \leq \binom{p+1}{2} - s \Leftrightarrow 4s \leq p^2 - 1.$$

Because of our hypotheses (14) we also conclude that $d_p \leq d$.

Therefore, $d_p \leq d \leq D_p$ and, by Proposition 2.2 in [1], there exists a smooth connected curve C of degree $d = \binom{p+1}{2} - s$ such that

$$H^0(\mathbb{P}^3, \mathcal{I}_C(p-1)) = H^1(\mathbb{P}^3, \mathcal{I}_C(p-1)) = H^2(\mathbb{P}^3, \mathcal{I}_C(p-1)) = 0.$$

In addition, by Corollary 2.4 and Remark 2.6 in [1], this curve is not contained in any surface of degree less than p , it has maximal rank and arithmetic genus

$$p_a = (p-1)d + 1 - \binom{p+2}{3} = \frac{(2p+3)(p-1)(p-2)}{6} - s(p-1).$$

Now, we may apply Corollary 5.1 in [6] to conclude that this curve C has a p -linear resolution, that is, the ideal sheaf \mathcal{I}_C has a p -linear resolution, and that this resolution is of the following type

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2-p)^s \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1-p)^{2s+p} \rightarrow \mathcal{O}_{\mathbb{P}^3}(-p)^{s+p+1} \rightarrow \mathcal{I}_C \rightarrow 0.$$

Finally, to prove that $H^2(\mathbb{P}^3, \mathcal{I}_C(p-4)) \neq 0$, consider the short exact sequence

$$0 \rightarrow \mathcal{I}_C(p-4) \rightarrow \mathcal{O}_{\mathbb{P}^3}(p-4) \rightarrow \mathcal{O}_C(p-4) \rightarrow 0,$$

from which we see that

$$H^2(\mathbb{P}^3, \mathcal{I}_C(p-4)) \cong H^1(C, \mathcal{O}_C(p-4)) \cong H^0(C, \omega_C(4-p)).$$

Since C is smooth, Proposition 5.2 in [7] guarantees the non-vanishing of $H^0(C, \omega_C(3-p))$. Then also $H^0(C, \omega_C(4-p)) \cong H^2(\mathbb{P}^3, \mathcal{I}_C(p-4)) \neq 0$, thus completing the proof. \square

We are now ready to construct, for each solution of (5), a vector bundle on the 3-dimensional projective space with the required properties.

Suppose (a, b, c) is a solution of (5), and hence, as stated in Proposition 2.6, of the form (12). Let

$$p = b - 2a = -\alpha\beta - 2\gamma\delta - 2(\beta^2 + 2\delta^2)$$

and

$$s = a = \frac{\alpha^2 + 2\gamma^2 - 2\beta^2 - 4\delta^2 + 2\alpha\beta + 4\gamma\delta}{8}.$$

Observe that p is even for α is even. If p and s are positive integers and furthermore $(p - 2s)^2 \geq 4s$, we know from Proposition 3.2 that there is a smooth curve C whose ideal sheaf has a linear resolution of the form

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2 - p)^s \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1 - p)^{2s+p} \rightarrow \mathcal{O}_{\mathbb{P}^3}(-p)^{s+p+1} \rightarrow \mathcal{I}_C \rightarrow 0.$$

Twisting this sequence by $\mathcal{O}_{\mathbb{P}^3}(p)$ and rewriting it in terms of a , b and c , we get

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2)^a \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^b \rightarrow \mathcal{O}_{\mathbb{P}^3}^{b-a+1} \rightarrow \mathcal{I}_C(b - 2a) \rightarrow 0.$$

Moreover, also from Proposition 3.2 we know that $H^2(\mathbb{P}^3, \mathcal{I}_C(b - 2a - 4)) \neq 0$. Since $H^2(\mathbb{P}^3, \mathcal{I}_C(b - 2a - 4)) \cong \text{Ext}^1(\mathcal{I}_C(b - 2a), \mathcal{O}_{\mathbb{P}^3})$, we can take a non-trivial element $e \in \text{Ext}^1(\mathcal{I}_C(b - 2a), \mathcal{O}_{\mathbb{P}^3})$. Then we have a diagram

$$\begin{array}{ccccccc}
& & & & 0 & & \\
& & & & \downarrow & & \\
& & & & \mathcal{O}_{\mathbb{P}^3}(-2)^a & & \\
& & & & \downarrow & & \\
& & & & \mathcal{O}_{\mathbb{P}^3}(-1)^b & & \\
& & & & \downarrow & & \\
0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^{c-b+a-1} & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^c & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^{b-a+1} \longrightarrow 0 \\
& & & & \searrow \phi & & \downarrow \\
e : & 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^{c-b+a-1} & \longrightarrow & E & \longrightarrow \mathcal{I}_C(b - 2a) \longrightarrow 0 \\
& & & & & & \downarrow \\
& & & & & & 0
\end{array}$$

where E is a representative of the corresponding equivalence class of e . Applying the covariant functor $\text{Hom}(\mathcal{O}_{\mathbb{P}^3}^c, -)$ to the bottom row we get the short exact sequence

$$0 \rightarrow \text{Hom}(\mathcal{O}_{\mathbb{P}^3}^c, \mathcal{O}_{\mathbb{P}^3}^{c-b+a-1}) \rightarrow \text{Hom}(\mathcal{O}_{\mathbb{P}^3}^c, E) \rightarrow \text{Hom}(\mathcal{O}_{\mathbb{P}^3}^c, \mathcal{I}_C(b - 2a)) \rightarrow 0.$$

Then, taking the morphism $\phi \in \text{Hom}(\mathcal{O}_{\mathbb{P}^3}^c, \mathcal{I}_C(b - 2a))$ there is a non-trivial morphism $\varphi \in \text{Hom}(\mathcal{O}_{\mathbb{P}^3}^c, E)$ that allows us to complete the above diagram

to the following commutative diagram

$$\begin{array}{ccccccc}
& & & 0 & & & 0 \\
& & & \downarrow & & & \downarrow \\
& & & \mathcal{O}_{\mathbb{P}^3}(-2)^a & \equiv & \mathcal{O}_{\mathbb{P}^3}(-2)^a & \\
& & & \downarrow & & & \downarrow \\
& & & \mathcal{O}_{\mathbb{P}^3}(-1)^b & \equiv & \mathcal{O}_{\mathbb{P}^3}(-1)^b & \\
& & & \downarrow & & & \downarrow \\
0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^{c-b+a-1} & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^c & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^{b-a+1} \longrightarrow 0 \\
& & & & \searrow \phi & & \downarrow \\
0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^{c-b+a-1} & \longrightarrow & E & \longrightarrow & \mathcal{I}_C(b-2a) \longrightarrow 0 \\
& & & & \downarrow & & \downarrow \\
& & & & 0 & & 0
\end{array}$$

The left column is a resolution of E and hence $\text{hd}(E) \leq 2$. To check that the resolution is minimal, that is the homological dimension of E is exactly 2, observe that one can easily deduce from the diagram that $H^1(E(-2)) \neq 0$. But this would contradict Lemma 3.1 in [11] if we had $\text{hd}(E) \leq 1$.

We have just proved our main theorem.

Theorem 3.3. *Suppose (a, b, c) is a solution of (5). If $a > 0$, $b - 2a > 0$ and $(b - 2a)^2 \geq 4a$, then there is a vector bundle E of homological dimension 2 and linear resolution*

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2)^a \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^b \rightarrow \mathcal{O}_{\mathbb{P}^3}^c \rightarrow E \rightarrow 0.$$

In particular, E satisfies $\chi(\text{End } E) = 1$.

We can apply Theorem 3.3 to get infinitely many families of such vector bundles. For example, consider the solutions of (5) with $\beta = \delta = 1$ and $\alpha = \gamma - 1$ even. In this case, we have

$$\begin{aligned}
a &= \frac{(\gamma - 1)^2 + 2\gamma^2 - 6 + 2(\gamma - 1) + 4\gamma}{8} = \frac{3\gamma^2 + 4\gamma - 7}{8}, \\
b &= \frac{(\gamma - 1)^2 + 2\gamma^2 - 18}{4} = \frac{3\gamma^2 - 2\gamma - 17}{4}, \\
c &= \frac{(\gamma - 1)^2 + 2\gamma^2 - 6 - 2(\gamma - 1) - 4\gamma}{8} = \frac{3\gamma^2 - 8\gamma - 3}{8},
\end{aligned}$$

where γ is an odd integer. Moreover, conditions

$$\begin{cases} a > 0 & \Leftrightarrow 3\gamma^2 + 4\gamma - 7 > 0 \\ b > 2a & \Leftrightarrow \gamma < -5/3 \\ (b - 2a)^2 \geq 4a & \Leftrightarrow 3\gamma^2 + 22\gamma + 39 \geq 0 \end{cases}$$

imply γ is an odd integer less than or equal to -5 . Hence, by Theorem 3.3, for each odd integer $\gamma \leq -5$ there is a vector bundle E_γ of rank 3 and homological dimension 2 defined by

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2)^{\frac{3\gamma^2+8\gamma-3}{8}} \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^{\frac{3\gamma^2+2\gamma-17}{4}} \rightarrow \mathcal{O}_{\mathbb{P}^3}^{\frac{3\gamma^2-4\gamma-7}{8}} \rightarrow E_\gamma \rightarrow 0.$$

For instance, if $\gamma = -5$, we get the following rank 3 vector bundle of homological dimension 2 with resolution

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2)^4 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^{12} \rightarrow \mathcal{O}_{\mathbb{P}^3}^{11} \rightarrow E_{-5} \rightarrow 0$$

satisfying $\chi(\text{End } E_{-5}) = 1$.

More generally, we can state the existence of the following family of rank 3 vector bundles on \mathbb{P}^3 .

Corollary 3.4. *For any odd integer $\gamma \leq -5$, there is a rank 3 vector bundle E_γ defined by a minimal linear resolution of type*

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2)^{\frac{3\gamma^2+4\gamma-7}{8}} \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^{\frac{3\gamma^2-2\gamma-17}{4}} \rightarrow \mathcal{O}_{\mathbb{P}^3}^{\frac{3\gamma^2-8\gamma-3}{8}} \rightarrow E_\gamma \rightarrow 0.$$

In particular, E satisfies $\chi(\text{End } E) = 1$.

In Proposition 4.3 in [11] the authors had proved the existence of vector bundles of homological dimension 2 and linear resolution

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2)^{u_s} \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^{u_{s+1}} \rightarrow \mathcal{O}_{\mathbb{P}^3}^{4u_{s+1}-10u_s} \rightarrow E \rightarrow 0, \quad (15)$$

and satisfying $\chi(\text{End } E) = 1$, where $\{u_s\}_{s \geq 0}$ is the sequence

$$\begin{cases} u_0 = 0 \\ u_1 = 1 \\ u_{s+2} = 4u_{s+1} - u_s \end{cases}$$

Hence, the 3-uples of the form $(a, b, c) = (u_s, u_{s+1}, 4u_{s+1} - 10u_s)$ are solutions of the Diophantine equation (5). It is obvious that these solutions form a subset of the set of general solutions obtained in this work. So a natural

question is how to determine when a general solution (12) comes from a solution of this form.

We can also ask whether Theorem 3.3 gives us all vector bundles, up to isomorphism, of this type (in the case of homological dimension 1 we have a description of all vector bundles with linear resolution of length 1 and $\chi(\text{End } E) = 1$).

In addition, note that the hypotheses $a > 0$ and $b - 2a > 0$ in the theorem are necessary restrictions on a and b in order to E to exist. Obviously, if E exists then one must have $a > 0$. In general, among all solutions of (5) the only feasible ones are those 3-uples (a, b, c) such that

$$\begin{aligned} a > 0 &\Leftrightarrow (\alpha + \beta)^2 + 2(\gamma + \delta)^2 > 3(\beta^2 + 2\delta^2) \\ b > 0 &\Leftrightarrow \alpha^2 + 2\delta^2 > 6(\beta^2 + 2\delta^2) \\ c > 0 &\Leftrightarrow (\alpha - \beta)^2 + 2(\gamma - \delta)^2 > 3(\beta^2 + 2\delta^2) \\ a \leq b &\Leftrightarrow (\alpha - \beta)^2 + 2(\gamma - \delta)^2 \geq 11(\beta^2 + 2\delta^2) \end{aligned}$$

But we immediately observe that the fourth inequality implies the third one. And it is also easy to check that if the first and the last conditions hold then the second one also holds. So, we need only to consider the first and the fourth inequalities, $a > 0$ and $a \leq b$. Proposition 3.1 implies that the feasible solutions can be further restricted to the two conditions in the theorem, $a > 0$ and $b - 2a > 0$.

Acknowledgements

Sérgio Mendes was partially supported by FCT through CMA-UBI (project UID/MAT/00212/2019). Rosa M. Miró-Roig was partially supported by MTM2016-78623-P. Helena Soares was partially supported by CIMA – Centro de Investigação em Matemática e Aplicações, Universidade de Évora, project UIDB/04674/2020 (Fundação para a Ciência e Tecnologia).

References

- [1] E. Ballico, G. Bolondi, P. Ellia and R. M. Miró-Roig, Curves of maximum genus in range A and stick-figures, *Trans. Amer. Math. Soc.* 349 (1997), 4589–4608.
- [2] G. Bohnhorst and H. Spindler, The stability of certain vector bundles on $\mathcal{O}_{\mathbb{P}^n}$, *Complex algebraic varieties (Bayreuth, 1990)*, 39–50, *Lecture Notes Math.*, 1507, Springer, Berlin, 1992.

- [3] M. C. Brambilla, Cokernel bundles and Fibonacci bundles, *Math. Nachr.* 281 (2008), 499–516.
- [4] J. W. S. Cassels, *Rational Quadratic Forms*, Dover, New York, 1978. (1994) 241–291.
- [5] I. Dolgachev and M. Kapranov, Arrangements of hyperplanes and vector bundles on \mathbb{P}^n , *Duke Math. J.* 71 (1984), 633–664.
- [6] D. Eisenbud and S. Goto, Linear free resolutions and minimal multiplicity, *J. Algebra* 88 (1984), 89–133.
- [7] R. Hartshorne, R. M. Miró-Roig, On the intersection of ACM curves in \mathbb{P}^3 , *J. Pure Appl. Algebra* 219 (2015), 3195–3213.
- [8] K. Ireland, M. Rosen, *A Classical Introduction to Modern Number Theory*, Springer, New York, 1990.
- [9] M. Jardim, R. V. Martins, Linear and Steiner Bundles on Projective Varieties, *Commun. Algebra* 38 (2010), 2249–2270.
- [10] S. Marchesi, D. M. Prata, Simplicity and exceptionality of syzygy bundles over \mathbb{P}^n . *Annali di Matematica* 195 (2016), 41–58.
- [11] R. M. Miró-Roig and H. Soares, Exceptional bundles of homological dimension k , *Forum Math.* 29 (2017), 701–715.
- [12] J.-P. Serre, *Course d’Arithmétique*, Coll. SUP: Le Mathématicien, Vol. 2, Presses Universitaires de France, Paris, 1970.
- [13] H. Soares, *Steiner Vector Bundles on Algebraic Varieties*, Ph.D. thesis, University of Barcelona, 2008.
- [14] A. Weil, *Number Theory: An Approach Through History*, Birkhäuser, Boston, Basel and Stuttgart, 1984.

S. Mendes, ISCTE, Av. das Forças Armadas, 1649-026, Lisbon, Portugal
 Email: sergio.mendes@iscte.pt

R. M. Miró-Roig, Gran Via de les Corts Catalanes, Barcelona, Spain
 Email: miro@ub.edu

H. Soares, ISCTE, Av. das Forças Armadas, 1649-026, Lisbon, Portugal
 Email: helena.soares@iscte.pt